

Topographic Effects on Ocean Mixing

Chris Garrett
Department of Physics and Astronomy
Elliott Building
University of Victoria
P.O. Box 3055 Victoria, BC
Canada, V8W 3P6
phone: 250-721-7702 fax: 250-721-7715 email: garrett@uvphys.phys.uvic.ca

Award # N00014-9310479
<http://maelstrom.seos.uvic.ca/>

LONG-TERM GOALS

To understand and parameterize interior and near-boundary mixing processes.

To understand the dynamical processes occurring in the surface layer of the ocean and parameterize them in ocean models.

To understand the physical oceanography of straits and semi-enclosed seas.

OBJECTIVES

An ongoing objective is to exploit Juan de Fuca Strait as a natural laboratory for the study of rotating stratified shear flows with sloping lateral boundaries. In particular I would like to understand and quantify vertical and lateral transfer of momentum and scalars, the causes and role of cross-strait secondary circulation, and the comparative importance, magnitude and parameterization of internal and near-boundary mixing.

In the surface mixed layer I would like to understand and parameterize the effects of various processes responsible for mixing and for air-sea gas exchange.

For straits I seek to determine the relative importance of hydraulic and frictional processes, and to understand the nature and causes of variations on a variety of time scales.

An ongoing study also seeks to elucidate the comparative importance of turbulence and zooplankton for acoustic backscatter at various frequencies.

APPROACH

For the last several summers we have conducted observational studies in Juan de Fuca Strait involving one or more bottom-mounted 300 kHz broadband ADCPs, temperature and conductivity moorings, and CTD profiles and “tow-yos”. Senior Research Associate Richard Dewey assumes much of the responsibility for this, with assistance from postdoctoral fellows (particularly Kate Stansfield) and graduate students Michael Ott, Keir Colbo, Tetjana Ross and Steven Stringer.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2001		2. REPORT TYPE		3. DATES COVERED 00-00-2001 to 00-00-2001	
4. TITLE AND SUBTITLE Topographic Effects on Ocean Mixing				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Physics and Astronomy,,Elliott Building,University of Victoria,P.O. Box 3055 Victoria, BC,,Canada, V8W 3P6, ,				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 7	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			




Figure 2. Internal tide energy flux spectra as a function of the horizontal wave number (and equivalent vertical mode number) for (a) the East Pacific Rise and (b) the Mid-Atlantic Ridge. The spectra are red, showing that most of the flux goes into low modes. The wavenumber equal to the tidal excursion is marked and is approximately 0.01 radian/m. The slope also becomes critical at about this scale for the Mid-Atlantic Ridge, but is still subcritical for the East Pacific Rise.

IMPACT/APPLICATIONS

Our results on internal Reynolds stresses in stratified flows have implications for the modeling of internal friction in estuarine and other models, and show the importance of direct measurements rather than relying on circumstantial evidence for verification of model parameterizations.

The role of internal tides in mixing the ocean is receiving much attention. Our study should help to clarify the importance and dominance of mixing remote from the generation site as well as providing accurate estimates of the energy flux from real topography.

The new results on hydraulic flows with entrainment could provide an intuitive basis for understanding real exchange flows.

The earlier results on bubble sizes will provide a useful input to models of air-sea gas exchange and acoustic propagation in the upper ocean.

Ongoing analysis of results from our Knight Inlet study will clarify the relative importance of turbulence and zooplankton in acoustic backscatter at various frequencies, and possibly provide clues to the effect of turbulence on zooplankton behaviour.

The proceedings of the 2001 'Aha Huliko'a, which I con-vened, should reach a wide audience interested in the interaction of stirring and mixing processes in the ocean.

TRANSITIONS

We are collaborating with Professor Parker MacCready and others at the University of Washington.

RELATED PROJECTS

The projects described above are also supported by Canadian funding agencies with equal or greater contributions to salaries and equipment and full provision of shiptime.

PUBLICATIONS

Garrett, C., M. Li and D. Farmer. 2000. The connection between bubble size spectra and energy dissipation rates in the upper ocean. *J. Phys. Oceanogr.*, **30**, 2163-2171.

Garrett, C. 2000. The dynamic ocean. In "*Perspectives in Fluid Dynamics*", Cambridge University Press, 507-556.

Ross, T., C. Garrett and P.-Y. Le Traon. 2000. Western Mediterranean sea level rise: changing exchange flow through the Strait of Gibraltar. *Geophys. Res. Lett.*, **27**, 2949-2952.

Garrett, C. 2001. An isopycnal view of near-boundary mixing and associated flows. *J. Phys. Oceanogr.*, **31**, 138-142.

Garrett, C. 2001. What is the "near-inertial" band and why is it different from the rest of the internal wave spectrum? *J. Phys. Oceanogr.*, **31**, 962-971.

Garrett, C. 2001. Internal waves. *Encyclopedia of Ocean Sciences*.

Stansfield, K., C. Garrett and R. Dewey. 2001. Calculating Thorpe scales and vertical mixing from CTD data, with application to Juan de Fuca Strait. *J. Phys. Oceanogr.*, (in press).

Garrett, C. 2001. Stirring and mixing: What are the rate-controlling processes? *Proceedings of the Twelfth 'Aha Huliko'a Hawaiian Winter Workshop*, (in press).

Gerdes, F., C. Garrett and D. Farmer. 2001. A note on hydraulics with entrainment. *J. Phys. Oceanogr.*, (in press).

Garrett, C. and L. St. Laurent. 2001. Aspects of deep ocean mixing. *J. Oceanogr.*, (in press).

St. Laurent, L. and C. Garrett. 2001. The role of internal tides in mixing the deep ocean. *J. Phys. Oceanogr.*, (submitted).

Ott, M. W., R. Dewey and C. Garrett. 2001. Reynolds stresses and secondary circulation in a stratified rotating shear flow. *J. Phys. Oceanogr.*, (submitted).

Müller, P. and C. Garrett. 2001. From stirring to mixing in a stratified ocean. *Oceanography*, (submitted).

Garrett, C. 2001. Tidal power generation by continuous flow. *Can. Appl. Math. Quarterly*, (submitted).